

REFERENCES

1. Aleksandrova N.V. *Matematicheskiye terminy* [Mathematical terms]. Moscow: Vysshaya shkola, 1978. 190 p. (In Russian)
2. Bronstein I.I., Semendyaev K.A. *Spravochnik po matematike* [Handbook of mathematics]. M.: Nauka, 1980. 975 p. (In Russian)
3. Fikhtengolts G.M. *Kurs differentsial'nogo i integral'nogo ischisleniya* [Course of differential and integral calculus]. Vol. 2. M.: Nauka, 1969. 800 p. (In Russian)
4. Yanke E., Emde F., Lesh F. *Special functions*. M.: Nauka, 1977. 342 p. (In Russian)
5. Milne-Thomson L. *Elliptic integrals. Reference book on special functions*. Eds. Abramowitz M., Stegun I. M.: Nauka, 1979. Pp. 401–441.
6. Anakhaev K.N. On methods of calculating potential (seepage) flows on the basis of Jacobi elliptic integrals. *Power Technology and Engineering*. 2008. Vol. 42. No. 5. Pp. 273–276. DOI: 10.1007/s10749-009-0052-0
7. Anakhaev K.N. Complete elliptic integrals of the third kind in problems of mechanics. *Doklady Physics*. 2017. Vol. 62. No. 3. Pp. 133–135. DOI: 10.1134/S1028335817030053
8. Anakhaev K.N. Elliptic integrals in nonlinear problems of mechanics. *Doklady Physics. Physics Reports* 2020. Vol. 65. No. 4. Pp. 142–146. DOI: 10.1134/S1028335820040011
MSC: 33, 45

LENGTH OF LEMNISCATA BERNOULLI ARC

K.N. ANAKHAEV

Institute of Applied Mathematics and Automation –
branch of Kabardino-Balkarian Scientific Center of the Russian Academy of Sciences
360000, Russia, Nalchik, 89 A Shortanov street

Annotation. The Bernoulli lemniscate curve can be used in applied studies of the shape of sliding and the shape of geophysical masses and natural objects, conjugations of trajectories of straight and rounded paths, movements of rod-hinge structures, equal gravitational movements along the lemniscate curve and its chord (tautochronicity), etc. The exact determination of the length of the lemniscate arc is due to the need to use an incomplete elliptical integral of the 1st kind (ungraspable), which makes it difficult to carry out analytical calculations, etc. The proposed elementary dependence for determining the length of the Bernoulli lemniscate arc gives a very close coincidence ($< 1-2\%$) with the basic data of the numerical solution and is recommended for use in solving applied problems in various fields of science and technology.

Keywords: 4-order curves, Bernoulli lemniscate, length of lemniscate arc, elliptic integrals of 1 genus, tautochronicity of curve

Information about the author

Anakhaev Koshkinbai Nazirovich, Doctor of Technical Sciences, Professor, Chief Researcher of the Department of Mathematical Modeling of Geophysical Processes, Institute of Applied Mathematics and Automation – branch of Kabardino-Balkarian Scientific Center of the Russian Academy of Sciences;

360000, Russia, Nalchik, 89 A Shortanov street;

anaha13@mail.ru, ORCID: <https://orcid.org/0000-0003-4357-4349>

